

An Evaluation of Potential Repellents for Botta's Pocket Gopher

Gary W. Witmer and Rachael S. Moulton

USDA APHIS WS National Wildlife Research Center, Fort Collins, Colorado

Roger A. Baldwin

Dept. of Wildlife, Fish, and Conservation Biology, University of California, Davis, California

ABSTRACT: Pocket gophers are fossorial rodents that cause substantial damage to crops, reforestation, and property. We tested potential repellents to identify candidates to reduce irrigation tubing damage. We dipped carrot chunks in the test materials, using mineral oil as the solvent. Gophers prefer tuberous roots and, when kept in captivity, are often fed carrots as part of their maintenance diet. None of the materials tested proved effective as repellents, even at concentrations as high as 20% active ingredient or in combinations. Wood blocks soaked in a few of the repellents received nearly significantly less damage than the control blocks and could be looked at further as repellents. However, it appears that the quest for an effective gopher repellent will continue to elude researchers.

KEY WORDS: Botta's pocket gopher, drip irrigation, pocket gopher, repellent, *Thomomys bottae*, wildlife damage

Proc. 27th Vertebr. Pest Conf. (R. M. Timm and R. A. Baldwin, Eds.)
Published at Univ. of Calif., Davis. 2016. Pp. 325-331.

INTRODUCTION

There are numerous species of pocket gophers in North America with most species belonging to the genera *Geomys* and *Thomomys* (Nowak 1991). Pocket gophers cause various types of damage to agricultural and rangeland resources and to reforestation (Witmer and Engeman 2007). Pocket gophers (*Thomomys* spp.) are generally considered one of the most damaging wildlife pests in California (Marsh 1992, Clark 1994). A recent study estimated average losses ranging from 5.3-8.8% across a variety of crops in CA (Baldwin et al. 2014), with one study showing a loss of 36.5% of annual production in alfalfa in fields with high density gopher populations (Smallwood and Geng 1997). The most widespread pocket gopher in California is the Botta's pocket gopher, *Thomomys bottae* (Case and Jasch 1994).

Primary control options for pocket gophers include trapping, burrow fumigation with aluminum phosphide, and baiting with rodenticides (Barnes et al. 1982, Baroch and Poche 1985, Evans et al. 1990, Case and Jasch 1994, Witmer and Engeman 2007, Baldwin 2012, Baldwin et al. 2014). Both trapping and burrow fumigation can be highly effective at controlling pocket gophers (Lewis and O'Brien 1990, Proulx 1997, Baker 2004), but are typically more time consuming and costly than baiting (Marsh 1992, Engeman and Witmer 2000). As such, baiting is often preferred by many growers, pest control advisors, and pest control operators. Three baits are used to control pocket gophers: strychnine, zinc phosphide, and first generation anticoagulants.

Pocket gophers are also damaging sub-surface drip irrigation tubes in agricultural fields in California. Repellents might provide a measure of protection to this tubing, if an effective repellent can be identified. The material would need to be added to the irrigation water or used as a coating on the tubing. Past efforts to identify effective gopher repellents have generally shown little promise (Witmer et al. 1997, Witmer et al. 1998) with a few exceptions, such as predator odors (Lindgren et al.

1997, Witmer et al. 1997) and capsaicin (Shumake et al. 1999, Sterner et al. 2005). An irrigation system company, Netafim Ltd., (Fresno, CA) requested us to revisit this situation and conduct cage trials to identify some effective repellents that they could then test in the field. Colleagues in Germany, working on vole damage, have identified some potential candidates that they found to be effective repellents for voles. We tested a few additional materials, based on the results of published scientific literature (e.g., papers in Mason 1997). A number of these materials are used as bird repellents, a few are used as mammal repellents, and some are used as insect repellents. Many are "essential oils" derived from plants and used in aromatic and other therapies. However, none are registered as pocket gopher repellents. The materials were considered to be mild irritants and potentially cause short-term irritation to the skin, eyes, gastrointestinal tract, mouth, and respiratory tract.

The objective of this study was to identify effective repellents to reduce damage by pocket gophers. We determined the efficacy of the potential repellents on wild-caught Botta's pocket gophers in a climate-controlled animal room of the USDA National Wildlife Research Center (NWRC) in Fort Collins, CO. We hypothesized that one or more of these materials would exhibit a high level of irritation to the olfactory, taste, or trigeminal receptors. We further hypothesized that some of the test materials would exhibit a high repellency, as determined by the amount of treated preferred foods consumed by gophers, using carrots as the preferred food item. We also monitored the amount of chewing done by gophers on treated wood blocks.

METHODS

Pocket gophers (henceforth, "gophers") used in this study were Botta's pocket gophers live-trapped in California and transported to the NWRC, Fort Collins, CO. Gophers were kept in individual numbered shoebox cages in a climate-controlled animal room. The lights

were always off unless staff had to do tasks in the room, during which times the lights were manually turned on, but at only 25% normal intensity. The temperature was maintained at 70°F with ambient humidity (~10-30%). They were fed a maintenance diet of rodent chow pellets and carrot chunks and received water *ad libitum*. They were provided with bedding, a den tube, and material to chew on (wood chunks). There was a 2-week quarantine period before the study began. Gophers were randomly assigned to treatment and control groups with three gophers per group. Each group contained at least one male or one female. The group size was kept small so that we could test more materials. However, for one treatment group (ProTec-T) we used five gophers because the study sponsor wanted a good test of their product. The weight, sex, cage number, and treatment of each gopher were recorded before the initiation of a trial. Gophers continued to receive the maintenance diet throughout the trial. On day one of the trial, one piece of carrot (each about 10-15 g) was added to each cage after it had been dipped in the repellent test material and weighed. Solutions of repellent test materials were prepared at the specified concentration by adding the chemical to mineral oil (the solvent used). Control gophers were maintained on the maintenance diet only throughout the trials. Prior to dipping, a piece of colored toothpick was stuck through the center of each treated carrot piece so that the remaining treated pieces could be identified from any remaining untreated pieces. One day later, any remaining treated carrot pieces were taken from the cage and their weight recorded before being returned to the cage for another full day. At the end of the 2-day exposure period, any remaining treated carrot pieces were removed, weighed, and disposed of. The concentration of repellent solutions was increased in each subsequent trial (Table 1). We also tried a few combinations of repellent materials (Table 1).

We conducted a final trial using gophers that had not been used before, but had shown a propensity to chew on the wood blocks. Wood blocks were soaked in a candidate repellent solution for 24 hours. Then each block was weighed before being placed into one of the gopher cages. As before, there were three gophers per treatment group. Three gophers were in a control group, which received a wood block that had been soaked only in mineral oil. Three days later, the wood blocks were removed and weighed so that the total amount removed (in grams) by gopher chewing could be determined. Gophers were examined daily and notes on their condition were recorded. The Attending Veterinarian or an Animal Care staff was contacted if any gopher appeared to be in more than momentary pain or distress. All surviving gophers were euthanized at the end of the study.

T-tests and ANOVA tests were used on some of the data sets to determine if a significant difference existed in consumption of treated carrots versus control carrots, and between treated and untreated wood blocks.

RESULTS AND DISCUSSION

A sizable number of potential repellents were tested in our study. Some of these are already known to have

some repellency with certain vertebrate or invertebrate species. Additionally, some have been identified in research studies as having repellency to some rodent species (e.g., Fischer et al. 2013, Cowan et al. 2014). Unfortunately, we did not find any of the tested materials to show significant repellency during our trials with Botta's pocket gopher. This was true even when the concentration was increased and when we combined two or three of the active ingredients (see Tables 1 and 2). We started with relatively low concentrations (generally 5% active ingredient), because we assumed only a low concentration in crop irrigation water would be acceptable so as to not affect food crop flavor or other attributes. One test material appeared to show promise as a repellent after one day of feeding on the dipped carrots. This was 10% 2-undecanone (also known as methyl nonyl ketone). Only 53% of the carrots were eaten by the gophers versus 84% of the control carrots ($T = 2.18$, $P = 0.095$). However, after the second day, the differences in the percentages eaten was much less, at 80% and 100%, respectively ($T = 1.58$, $P = 0.190$). One material tested, 5% bergamot oil, almost appeared to be an attractant, with 99% of the dipped carrots eaten after one day versus the 77% of the control carrots. However, the difference was not significant ($T = 1.75$, $P = 0.156$).

In the second part of the study, we used wood blocks that had been soaked in the potential repellent materials (Table 2). Three of the materials tested had nearly significantly less gnawing of the wood blocks than the control blocks ($F = 3.82$, $P = 0.057$). These materials were 10% Pulegone, 10% 2-undecanone, and the combination of these active ingredients: 10% 2-undecanone, 10% black pepper oil, and 10% methyl anthranilate. These materials could be further researched for their potential as gopher repellents.

Materials that have repellency for a particular rodent species may not show repellency for other rodent species. For example, Fischer et al. (2013) found methyl nonyl ketone (= 2-undecanone) to be effective with common voles (*Microtus arvalis*) in Europe, while we did not find it to be effective with prairie voles (*M. ochrogaster*) in North America (Witmer et al. 2000). Of course, regional difference can also exist in how the same species of rodent responds to an odor or a toxicant.

We also tested anthraquinone, which is a registered bird repellent. It is being looked at as a potential addition to rodenticides to reduce the consumption of rodenticide baits by non-target birds (Cowan et al. 2015). However, for rats, it acted as a rodent repellent even at very low concentrations (Cowan et al. 2015). In our study, however, it did not repel gophers even at a concentration of 25%.

In conclusion, it appears that the quest for an effective gopher repellent will continue to elude researchers. Other researchers have noted that potential repellents that work for some rodent species may not work for other rodent species, and that ones that work for one gender may not be very effective for the other gender (Hansen et al. 2015, Hansen et al. 2016). In our previous trials to identify effective gopher repellents, it was only predator odors that showed the most potential promise as repellents (Witmer et al. 1997). In their review, Lindgren et al.

Table 1. Carrot consumption after one day and two days by treatment groups and control groups of pocket gophers.

Group	Animal ID	Carrot Weight Day 0 (g)	Date Carrots Given	Carrot Weight Day 1 (g)	Amount Eaten Day 1 (g)	Percent Eaten Day 1	Carrot Weight Day 2 (g)	Amount Eaten Day 2 (g)	Percent Eaten Day 2	Total Amount Eaten (g)	Ave. (S.D.) Amount Eaten (g)	Total Percent Eaten	Ave. (S.D.) Percent Eaten (g)	
UND 5%	OB36F	10.16	11/5/14	4.66	5.50	54.13	1.11	3.55	34.94	9.05	9.67 (0.80)	89.07	96.36 (6.31)	
	OB39F	9.39	11/5/14	0.00	9.39	100.00	0.00	0.00	0.00	9.39		100.00		100.00
	OB53M	10.57	11/5/14	1.63	8.94	84.58	0.00	1.63	15.42	10.57		100.00		100.00
PUL 5%	OD21F	10.36	11/5/14	0.00	10.36	100.00	0.00	0.00	0.00	10.36	10.10 (1.09)	100.00	91.22 (11.91)	
	OB23M	11.46	11/5/14	8.82	2.64	23.04	2.56	6.26	54.62	8.90		77.66		96.00
	OB40F	11.50	11/5/14	3.55	7.95	69.13	0.46	3.09	26.87	11.04		100.00		100.00
PEP 5%	OB19F	11.72	11/5/14	0.00	11.72	100.00	0.00	0.00	0.00	11.72	10.51 (1.07)	100.00	100.00 (0.00)	
	OB21F	10.14	11/5/14	2.46	7.68	75.74	0.00	2.46	24.26	10.14		100.00		100.00
	OB29M	9.68	11/5/14	0.00	9.68	100.00	0.00	0.00	0.00	9.68		100.00		100.00
ANT 6%	OB12F	8.70	11/6/14	0.00	8.70	100.00	0.00	0.00	0.00	8.70	8.95 (0.43)	100.00	100.00 (0.00)	
	OB33F	9.45	11/6/14	0.00	9.45	100.00	0.00	0.00	0.00	9.45		100.00		100.00
	OB48M	8.71	11/6/14	0.00	8.71	100.00	0.00	0.00	0.00	8.71		100.00		100.00
CIN 5%	OB11F	8.62	11/5/14	0.00	8.62	100.00	0.00	0.00	0.00	8.62	8.60 (0.66)	100.00	100.00 (0.00)	
	OB16F	7.93	11/5/14	0.00	7.93	100.00	0.00	0.00	0.00	7.93		100.00		100.00
	OD36AM	9.25	11/5/14	0.00	9.25	100.00	0.00	0.00	0.00	9.25		100.00		100.00
MET 5%	OB18F	8.46	11/5/14	6.10	2.36	27.90	0.00	6.10	72.10	8.46	8.77 (0.50)	100.00	100.00 (0.00)	
	OB20M	8.51	11/5/14	3.00	5.51	64.75	0.00	3.00	35.25	8.51		100.00		100.00
	OB22M	9.35	11/5/14	0.00	9.35	100.00	0.00	0.00	0.00	9.35		100.00		100.00
CON	OD28M	11.42	11/4/14	0.00	11.42	100.00	0.00	0.00	0.00	11.42	11.85 (0.56)	100.00	100.00 (0.00)	
	OD33F	11.66	11/4/14	3.05	8.61	73.84	0.00	3.05	26.16	11.66		100.00		100.00
	OB49F	12.48	11/4/14	5.47	7.01	56.17	0.00	5.47	43.83	12.48		100.00		100.00
BER 5%	OB05F	12.82	11/5/14	0.40	12.42	96.88	0.00	0.40	3.12	12.82	11.85 (0.85)	100.00	100.00 (0.00)	
	OB17M	11.48	11/5/14	0.00	11.48	100.00	0.00	0.00	0.00	11.48		100.00		100.00
	OB35F	11.24	11/5/14	0.00	11.24	100.00	0.00	0.00	0.00	11.24		100.00		100.00
UND 10%	OB36F	18.57	11/18/14	11.42	7.15	38.50	7.69	3.73	20.09	10.88	12.40 (2.53)	58.59	80.35 (20.79)	
	OB39F	15.32	11/18/14	4.55	10.77	70.30	0.00	4.55	29.70	15.32		100.00		100.00
	OB53M	13.34	11/18/14	6.44	6.90	51.72	2.34	4.10	30.73	11.00		82.46		82.46
PUL 10%	OD21F	12.70	11/18/14	2.74	9.96	78.43	0.00	2.74	21.57	12.70	13.67 (1.30)	100.00	100.00 (0.00)	
	OB23M	13.17	11/18/14	5.14	8.03	60.97	0.00	5.14	39.03	13.17		100.00		100.00

Group	Animal ID	Carrot Weight Day 0 (g)	Date Carrots Given	Carrot Weight Day 1 (g)	Amount Eaten Day 1 (g)	Percent Eaten Day 1	Carrot Weight Day 2 (g)	Amount Eaten Day 2 (g)	Percent Eaten Day 2	Total Amount Eaten	Ave. (S.D.) Amount Eaten (g)	Total Percent Eaten	Ave. (S.D.) Percent Eaten (g)
PEP 10%	OB40F	15.15	11/18/14	6.72	8.43	55.64	0.00	6.72	44.36	15.15		100.00	
	OB19F	13.14	11/18/14	0.52	12.62	96.04	0.00	0.52	3.96	13.14		100.00	
	OB21F	16.11	11/18/14	10.51	5.60	34.76	3.55	6.96	43.20	12.56	13.05 (0.45)	77.96	92.65 (12.72)
	OB29M	13.45	11/18/14	2.68	10.77	80.07	0.00	2.68	19.93	13.45		100.00	
ANT 12%	OB12F	11.99	11/18/14	0.00	11.99	100.00	0.00	0.00	0.00	11.99		100.00	
	OB33F	16.80	11/18/14	2.23	14.57	86.73	0.00	2.23	13.27	16.80	13.78 (2.63)	100.00	100.00 (0.00)
	OB48M	12.54	11/18/14	3.64	8.90	70.97	0.00	3.64	29.03	12.54		100.00	
	OB11F	14.83	11/18/14	0.00	14.83	100.00	0.00	0.00	0.00	14.83		100.00	
CIN 10%	OB16F	17.60	11/18/14	2.65	14.95	84.94	0.00	2.65	15.06	17.60	15.52 (1.84)	100.00	100.00 (0.00)
	OD36AM	14.12	11/18/14	0.80	13.32	94.33	0.00	0.80	5.67	14.12		100.00	
	OB18F	14.27	11/18/14	3.19	11.08	77.65	0.00	3.19	22.35	14.27		100.00	
	OB20M	14.29	11/18/14	0.00	14.29	100.00	0.00	0.00	0.00	14.29	13.61 (1.15)	100.00	97.99 (3.49)
MET 10%	OB22M	13.07	11/18/14	1.25	11.82	90.44	0.79	0.46	3.52	12.28		93.96	
	OD28M	14.47	11/18/14	0.00	14.47	100.00	0.00	0.00	0.00	14.47		100.00	
	OD33F	13.23	11/18/14	1.61	11.62	87.83	0.00	1.61	12.17	13.23	13.98 (0.66)	100.00	100.00 (0.00)
	OB49F	14.24	11/18/14	5.10	9.14	64.19	0.00	5.10	35.81	14.24		100.00	
BER 10%	OB05F	14.59	11/18/14	0.00	14.59	100.00	0.00	0.00	0.00	14.59		100.00	
	OB17M	14.29	11/18/14	0.00	14.29	100.00	0.00	0.00	0.00	14.29	13.95 (0.87)	100.00	100.00 (0.00)
	OB35F	12.96	11/18/14	0.00	12.96	100.00	0.00	0.00	0.00	12.96		100.00	
	OD01M	9.44	11/24/14	1.34	8.10	85.81	0.00	1.34	14.19	9.44		100.00	
Pro .5%	OD21AF	8.95	11/24/14	0.00	8.95	100.00	0.00	0.00	0.00	8.95		100.00	
	OD28M	10.30	11/24/14	0.00	10.30	100.00	0.00	0.00	0.00	10.30	10.19 (1.02)	100.00	100.00 (0.00)
	OD33F	10.76	11/24/14	4.52	6.24	57.99	0.00	4.52	42.01	10.76		100.00	
	OB49F	11.51	11/24/14	3.82	7.69	66.81	0.00	3.82	33.19	11.51		100.00	
UND 20%	OV2M	16.2	2/2/15	0.0	16.2	100.0	0.0	0.0	0.0	16.20		100.00	
	OV22M	10.4	2/2/15	0.0	10.4	100.0	0.0	0.0	0.0	10.40	12.83 (3.01)	100.00	100.00 (0.00)
	OV28F	11.9	2/2/15	0.0	11.9	100.0	0.0	0.0	0.0	11.90		100.00	
	OV12M	17.4	2/2/15	0.0	17.4	100.0	0.0	0.0	0.0	17.40		100.00	100.00 (0.00)
PUL 20%	OV18M	14.9	2/2/15	4.1	10.8	72.5	0.0	4.1	27.5	14.90	15.67 (1.50)	100.00	100.00 (0.00)

Group	Animal ID	Carrot Weight Day 0 (g)	Date Carrots Given	Carrot Weight Day 1 (g)	Amount Eaten Day 1 (g)	Percent Eaten Day 1	Carrot Weight Day 2 (g)	Amount Eaten Day 2 (g)	Percent Eaten Day 2	Total Amount Eaten (g)	Ave. (S.D.) Amount Eaten (g)	Total Percent Eaten	Ave. (S.D.) Percent Eaten (g)
Pro 1%	OV21F	14.7	2/2/15	9.3	5.4	36.7	3.0	9.3	63.3	14.70	10.67 (1.91)	100.00	
	OV6M	11.4	2/2/15	0.0	11.4	100.0	0.0	0.0	0.0	11.40		100.00	
	OV10M	12.7	2/2/15	4.2	8.5	66.9	0.0	0.0	0.0	8.50		66.93	88.98 (19.09)
	OV17F	12.1	2/2/15	0.0	12.1	100.0	0.0	0.0	0.0	12.10		100.00	
UBP 10%/10%	OV8M	7.9	2/2/15	0.0	7.9	100.0	0.0	0.0	0.0	7.90	9.33 (1.63)	100.00	
	OV15M	9.0	2/2/15	0.0	9.0	100.0	0.0	0.0	0.0	9.00		100.00	100.00 (0.00)
	OV16M	11.1	2/2/15	1.5	9.6	86.5	0.0	1.5	13.5	11.10		100.00	
	OV4M	13.3	2/2/15	0.0	13.3	100.0	0.0	0.0	0.0	13.30	11.77 (1.39)	100.00	100.00 (0.00)
UBPMA 10%/lea	OV13M	11.4	2/2/15	0.0	11.4	100.0	0.0	0.0	0.0	11.40		100.00	
	OV30F	10.6	2/2/15	1.5	9.1	85.8	0.0	1.5	14.2	10.60		100.00	
	OV9M	11.4	2/2/15	0.0	11.4	100.0	0.0	0.0	0.0	11.40		100.00	
	OV14M	16.2	2/2/15	3.1	13.1	80.9	0.0	3.1	19.1	16.20	12.53 (3.25)	100.00	100.00 (0.00)
CON	OV31F	10.0	2/2/15	0.0	10.0	100.0	0.0	0.0	0.0	10.00		100.00	
	OV03M	13.0	2/23/15	6.8	6.2	47.7	5.0	1.8	13.8	8.0	10.43 (2.40)	61.5	86.67 (21.78)
	OV15M	10.5	2/23/15	0.0	10.5	100.0	0.0	0.0	0.0	10.5		100.0	
	OV24F	13.0	2/23/15	5.9	7.1	54.6	0.2	5.7	43.8	12.8		98.5	
AVI 50%	OV01F	8.6	2/23/15	4.7	3.9	45.3	0.0	4.7	54.7	8.6	10.33 (1.51)	100.0	100.00 (0.00)
	OV11M	11.0	2/23/15	0.0	11.0	100.0	0.0	0.0	0.0	11.0		100.0	
	OV22M	11.4	2/23/15	0.0	11.4	100.0	0.0	0.0	0.0	11.4		100.0	
	OV20F	12.9	2/23/15	3.3	9.6	74.4	0.0	3.3	25.6	12.9	10.30 (2.25)	100.0	98.26 (3.01)
CON	OV27M	8.9	2/23/15	0.0	8.9	100.0	0.0	0.0	0.0	8.9		100.0	
	OV31F	9.6	2/23/15	2.1	7.5	78.1	0.5	1.6	16.7	9.1		94.8	
	OV15M	7.74	4/15/15	0.00	7.74	100.00	0.00	0.00	0.00	7.74	7.83 (0.17)	100.00	100.00 (0.00)
	OV20F	7.73	4/15/15	0.00	7.73	100.00	0.00	0.00	0.00	7.73		100.00	
Deer Off	OV31F	8.03	4/15/15	0.00	8.03	100.00	0.00	0.00	0.00	8.03		100.00	
	OV11M	10.39	4/15/15	0.00	10.39	100.00	0.00	0.00	0.00	10.39	8.87 (1.40)	100.00	100.00 (0.00)
	OV25F	7.62	4/15/15	0.00	7.62	100.00	0.00	0.00	0.00	7.62		100.00	
	OV30F	8.60	4/15/15	0.00	8.60	100.00	0.00	0.00	0.00	8.60		100.00	

Abbreviations: UND=2-undecanone; PUL=Pulegone; PEP=Black pepper oil; ANT=Anthraquinone; CIN=Cinnamamide; MET=Methyl anthranilate; CON=Control (mineral oil only); BER=Bergamot oil; PRO=Protect-T; UND=2-undecanone; PUL=Pulegone; PROT=Protect-T (undiluted); UBP=Undecanone (10%) and black pepper oil (10%); UBPMA=Undecanone (10%) and black pepper oil (10%); AVI=Avipel (undiluted from Scott Weimer, 50%); ANT=Anthranilate (25% in mineral oil); Deer Off=undiluted concentrate

Table 2. Amount of wood block removed by gopher chewing after three days by treatment groups and control group.

Group	Animal ID	Initial Block Weight(g)	Date Block Given	Final Block Weight(g)	Date Block Removed	Total Amount Chewed (g)	Ave. (S.D.) Amount Chewed (g)	Total Percent Chewed	Ave. (S.D.) Percent Chewed
UND 10%	OV2M	16.9	1/27/2015	15.7	1/30/2015	1.20	0.57 (0.57)	7.10	3.31 (3.39)
	OV22M	17.7	1/27/2015	17.3	1/30/2015	0.40		2.26	
	OV28F	17.2	1/27/2015	17.1	1/30/2015	0.10		0.58	
PUL 10%	OV12M	16.5	1/27/2015	16.2	1/30/2015	0.30	0.27 (0.15)	1.82	1.59 (0.91)
	OV18M	16.9	1/27/2015	16.8	1/30/2015	0.10		0.59	
	OV21F	16.9	1/27/2015	16.5	1/30/2015	0.40		2.37	
Prot-T	OV6M	19.2	1/27/2015	16.3	1/30/2015	2.90	2.63 (0.38)	15.10	13.21 (2.24)
	OV10M	20.5	1/27/2015	18.3	1/30/2015	2.20		10.73	
	OV17F	20.3	1/27/2015	17.5	1/30/2015	2.80		13.79	
UBP	OV8M	17.4	1/27/2015	17.0	1/30/2015	0.40	3.13 (4.65)	2.30	17.90 (26.56)
	OV15M	17.5	1/27/2015	9.0	1/30/2015	8.50		48.57	
	OV16F	17.6	1/27/2015	17.1	1/30/2015	0.50		2.84	
UBPMA	OV4M	16.3	1/27/2015	16.0	1/30/2015	0.30	0.80 (0.78)	1.84	4.69 (4.41)
	OV13M	17.4	1/27/2015	15.7	1/30/2015	1.70		9.77	
	OV30F	16.2	1/27/2015	15.8	1/30/2015	0.40		2.47	
CON	OV9M	17.2	1/27/2015	16.6	1/30/2015	0.60	2.87 (4.01)	3.49	15.97 (22.06)
	OV14M	16.8	1/27/2015	16.3	1/30/2015	0.50		2.98	
	OV31F	18.1	1/27/2015	10.6	1/30/2015	7.50		41.44	

Abbreviations and concentrations: UND=2-undecanone 10% (= methyl nonyl ketone); PUL=Pulegone 10%; Prot-T=Protect-T (undiluted; <1% propriety active ingredients); UBP=Undecanone (10%) and black pepper oil (10%); UBPMA=Undecanone (10%) and black pepper oil (10%) and methyl anthranilate (10%); CON=Control (mineral oil only)

(1997) also noted the potential for predator odors as repellent for various rodent species. However, in the current trials, even potential repellents that smelled like predator odors (sulfur-containing materials such as the Protect-T product) did not repel gophers. As suggested by Baldwin et al. (2014), additional research and development of effective methods will be necessary to reduce rodent damage to agriculture production.

ACKNOWLEDGMENTS

This study was conducted under the NWRC IACUC approved study protocol QA-2285. The study was funded by Netafim Irrigation, Inc., Fresno, California. We thank the landowners that allowed us to live trap pocket gophers on their properties in California. The mention of a commercial product or company does not represent an endorsement by the U.S. Government.

LITERATURE CITED

Baker, R. O. 2004. Field efficacy of Fumitoxin® (55% aluminum phosphide) tablets for controlling valley pocket gopher. *Proc. Vertebr. Pest Conf.* 21:253-257.
 Baldwin, R. A. 2012. The importance of aluminum phosphide for burrowing pest control in California. *Proc. Vertebr. Pest Conf.* 25:151-159.
 Baldwin, R. A. 2014. Determining and demonstrating the importance of training and experience for managing pocket gophers. *Wildl. Soc. Bull.* 38:628-633.

Baldwin, R. A., T. P. Salmon, R. H. Schmidt, and R. M. Timm. 2014. Perceived damage and areas of needed research for wildlife pests of agriculture. *Integrative Zoology* 9:265-279.
 Barnes, V. G., Jr., R. M. Anthony, J. Evans, and G. D. Lindsey. 1982. Evaluation of zinc phosphide bait for pocket gopher control on forest land. *Proc. Vertebr. Pest Conf.* 10:219-225.
 Baroch, J., and R. M. Poché. 1985. Preliminary field evaluation of a new formulation of Rozol (chlorophacinone) bait against pocket gophers in Colorado. *Proc. Gt. Plains Wildl. Damage Control Workshop* 7:138-144.
 Case, R. M., and B. A. Jasch. 1994. Pocket gophers. Pp. B17-B29 *in*: S. E. Hygnstrom, R. M. Timm, and G. E. Larson (Eds.), *Prevention and Control of Wildlife Damage*. Univ. of Nebraska Cooperative Extension, Lincoln, NE.
 Clark, J. P. (Editor). 1994. *Vertebrate Pest Control Handbook*, 4th Ed. California Department of Food and Agriculture, Sacramento, CA. 803 pp.
 Cowan, P., S. Brown, G. Forrester, L. Booth, and M. Crowell. 2015. Bird-repellent effects on bait efficacy for control of invasive mammal pests. *Pest Manage. Sci.* 71:1075-1081.
 Engeman, R. M., and G. W. Witmer. 2000. Integrated management tactics for predicting and alleviating pocket gopher (*Thomomys* spp.) damage to conifer reforestation plantings. *Integr. Pest Manage. Rev.* 5:41-55.

- Evans, J., G. H. Matschke, D. L. Campbell, P. L. Hegdal, and R. M. Engeman. 1990. Efficacy data for registration of strychnine grain baits to control pocket gophers (*Thomomys* spp.). *Proc. Vertebr. Pest Conf.* 14:82-86.
- Fischer, D., C. Imholt, A. Prokop, and J. Jacob. 2013. Efficacy of methyl nonyl ketone as an in-soil repellent for common voles (*Microtus arvalis*). *Pest Manage. Sci.* 69:431-436.
- Hansen, S. C., C. Stolter, and J. Jacob. 2015. The smell to repel: the effect of odors on the feeding behavior of female rodents. *Crop Protect.* 78:270-276.
- Hansen, S. C., C. Stolter, and J. Jacob. 2016. Effect of plant secondary metabolites on feeding behavior of microtine and arvicoline rodent species. *J. Pest Sci.* 89(4):955-963. doi: 10.1007/s10340-015-0723.
- Lewis, S. R., and J. M. O'Brien. 1990. Survey of rodent and rabbit damage to alfalfa hay in Nevada. *Proc. Vertebr. Pest Conf.* 14:116-117.
- Lindgen, P., T. Sullivan, and D. Crump. 1997. Review of synthetic predator odor semiochemicals as repellents for wildlife management in the Pacific Northwest. Pp. 217-230 *in*: R. Mason (Ed.), *Repellents in Wildlife Management: Proceedings of a Symposium*. USDA National Wildlife Research Center, Fort Collins, CO.
- Marsh, R. E. 1992. Reflections on current (1992) pocket gopher control in California. *Proc. Vertebr. Pest Conf.* 15:289-295.
- Mason, R. (Editor). 1997. *Repellents in Wildlife Management: Proceedings of a Symposium*. Proceedings of the Second DWRC Special Symposium (August 8-10, 1995, Denver, Colorado) National Wildlife Research Center, Fort Collins, CO.
- Nowak, R. 1991. *Walker's Mammals of the World*, 5th Ed., Vol. II. Johns Hopkins University Press, Baltimore, MD. 1,629 pp.
- Proulx, G. 1997. A northern pocket gopher (*Thomomys talpoides*) border control strategy: promising approach. *Crop Protect.* 16:279-284.
- Shumake, S., R. Sterner, and S. Gaddis. 1999. Repellents to reduce cable gnawing by northern pocket gophers. *J. Wildl. Manage.* 63:1344-1349.
- Smallwood, S., and S. Geng. 1997. Multiscale influences of gophers on alfalfa yield and quality. *Field Crops Res.* 49:159-168.
- Sterner, R., S. Shumake, S. Gaddis, and J. Bourassa. 2005. Capsaicin oleoresin: development of an in-soil repellent for pocket gophers. *Pest Manage. Sci.* 61:1202-1208.
- Witmer, G., and R. Engeman. 2007. Subterranean rodents as pests: the case of the pocket gopher. Pp. 285-299 *in*: S. Begall (Ed.), *Subterranean Rodents: News from Underground*. Springer-Verlag, Berlin.
- Witmer, G., R. Saylor, and M. Pipas. 1997. Repellent trials to reduce reforestation damage by pocket gophers, deer, and elk. Pp. 321-332 *in*: R. Mason (Ed.), *Repellents in Wildlife Management: Proceedings of a Symposium*. USDA National Wildlife Research Center, Fort Collins, CO.
- Witmer, G., M. Pipas, and J. Bucher. 1998. Field test of denatonium benzoate to reduce seedling damage by pocket gophers. *Crop Protect.* 17:35-39.
- Witmer, G., A. Hakim, and B. Moser. 2000. Investigations of methods to reduce damage by voles. *Proc. Wildl. Damage Manage. Conf.* 9:357-365.